



BANANA

APPLICATION SCHEDULE FOR BANANA\*\*

MONTHS AFTER PLANTATION	PRODUCTS	APPLICATION	METHOD AND DOSAGE
At the time of land preparation	AJAY AZO, AJAY POTASH and SPHURDA.	Biofix bio-fertilizers improves soil fertility naturally.	SOIL APPLICATION- Ajay Azo-2Kg + Ajay Potash-2Kg + Sphurda-2 Kg per acre
At the time of land preparation	AJAY NEEM PLUS	To improve efficiency of nitrogen providing fertilizers.	SOIL APPLICATION: Ajay Neem Plus-200 Kg/Acre.
	AJAY AZO, AJAY POTASH SPHURDA	Biofix bio-fertilizers improves soil fertility naturally.	DRIP /DRENCHING APPLICATION- Ajay Azo- 1 Litre + Ajay Potash- 1 Litre + Sphurda- 1 Litre per acre.
	AJAY VAM	For phosphorus solubilization, water absorption and root development.	SOIL APPLICATION: Apply 200 gm of AJAY VAM /Acre. Apply uniformly near the root zone of Plant.
	HUMIGUARD-PLUS	For white root development.	DRIP /DRENCHING APPLICATION- Humiguard Plus- 1 litre/Acre.
	BIO-FIGHTER	To preventive control of soil-borne larval pests in + primary stage.	DRIP APPLICATION / SOIL APPLICATION- Biofighter - 1 Kg/Acre in 200 litres of water.
3rd month	MICRO KOMBI-L	For balance nutrition of the crop.	DRIP / DRENCHING APPLICATION- Microkombi-L 500 ml/200 Litres of water per acre.
	BIOGUARD	For management of fungal disease and bacterial disease.	FOLIAR APPLICATION- Bioguard 2 ml/litre of water.
	BIOSAN	To minimise infestation of bacterial and insect pests.	FOLIAR APPLICATION- Biosan- 2 gm/litre of water.
	MITLAR	Biological control for sucking pests specially for Red Spider Mites.	FOLIAR APPLICATION- Mitar-1 to 1.5ml/Litre of water.
4th month	HUMIGUARD-PLUS	For white root development.	DRIP /DRENCHING APPLICATION- Humiguard Plus- 1 litre/Acre.
5th month	AJAY POTASH	Biofix bio-fertilizers improves soil fertility naturally.	DRIP /DRENCHING APPLICATION-Ajay Potash-1 Litre/Acre in 200 litres of water.
7th month	AJAY POTASH	Biofix bio-fertilizers improves soil fertility naturally.	DRIP /DRENCHING APPLICATION-Ajay Potash-1 Litre/Acre in 200 litres of water.
	MICRO KOMBI-L	For balance nutrition of the crop.	DRIP / DRENCHING APPLICATION- Microkombi-L 500 ml in 200 Litres of water per acre.
	KITOSTAR	Effective for management of Powdery mildew and Downy mildew.	FOLIAR APPLICATION- Kitostar-2ml/Litre of water.
Suitable to use in summer season	AJAY ANTI-STRESS	Helps in minimising the evaporation rate.	FOLIAR APPLICATION- Ajay Anti-Stress 1 to 2 ml/Litre of water,

- \* The above growth stages generally coincide with your plantation season. (State-wise plantation methods are different)
- \*\* The above given schedule taken as a Guideline for all stages of crop.
- Do not mix above Biological products with any chemical products. Spray during Early morning Or Late evening hours. Irrigation of Crops before 24-48 Hrs. of Bio-product applications is desirable.
- Keep minimum interval of 3 days between sprays of Biological products and Chemical products.

**Note:** The use of products of Ajay Biotech (India) Ltd. cannot be monitored by us in reality, we can only guarantee the quality of the products. Depending on the geographical area, spraying of fungicides and pesticides should be done according to and with the help of experts depending on crop disease, pest ratio, and climate change.