



ONION / GARLIC

APPLICATION SCHEDULE FOR ONION / GARLIC**

DAYS AFTER TRANSPLANTING	PRODUCTS	APPLICATION	METHOD AND DOSAGE
At the time of land preparation.	AJAY AZO, AJAY SPHURDA and AJAY POTASH,	To improve soil fertility naturally.	SOIL APPLICATION: Mix Ajay Azo- 2 Kg + Ajay Sphurda- 2 Kg + Ajay Potash- 2 Kg per acre .
	AJAY VAM and MICROKOMBI-F	For phosphorus solubilization, root development and for balance nutrition of the crop.	SOIL APPLICATION: Mix 100 gm of AJAY VAM and Microkombi 200 gm per acre.
1st Application 25 days after transplantation and 2nd Application 50 days after transplantation.	GROWRICH PLUS	To increase growth vigour of the crop.	FOLIAR APPLICATION- Growth Rich Plus- 1ml/litre of water.
	HUMIGUARD-PLUS	For white root development.	DRIP / DRENCHING APPLICATION- Humiguard-Plus 1 Litre per acre.
	BIO-GUARD	For management of fungal diseases and pests.	FOLIAR APPLICATION: Bio-guard- 1.5 to 2 ml/ litre of water.
	AJAY POTASH	To improve soil fertility naturally.	DRIP / DRENCHING APPLICATION: Ajay Potash- 1 Litre in 200 litres of water.
1st Application 25 days after transplantation and 2nd Application 50 days after transplantation.	BIOSAN	To minimise infestation of bacterial and insect pest	FOLIAR APPLICATION- Biosan- 2 gm/litre of water.
	MITLAR	Biological control for sucking pests.	FOLIAR APPLICATION- Mitlar-1 ml/litre of water.
	MICRO KOMBI-L	For balance nutrition of the crop.	DRIP / DRENCHING APPLICATION- Microkombi-L 500 ml in 200 ltr of water per acre.
	AJAY-SULF	For management of sulphur nutrition.	FOLIAR APPLICATION- Ajay Sulf-1 ml/litre of water.

- Repeat the sprays of dosages of Bioguard and Biosan, whenever required.
- * The above growth stages generally coincide with your plantation season. (State-wise plantation methods are different)
- ** The above given schedule taken as a Guideline for all stages of crop.
- Do not mix above Biological products with any chemical products. Spray during Early morning Or Late evening hours. Irrigation of Crops before 24-48 Hrs. of Bio-product applications is desirable.
- Keep minimum interval of 3 days between sprays of Biological products and Chemical products.

Note: The use of products of Ajay Biotech (India) Ltd. cannot be monitored by us in reality, we can only guarantee the quality of the products. Depending on the geographical area, spraying of fungicides and pesticides should be done according to and with the help of experts depending on crop disease, pest ratio, and climate change.