



TOMATO

APPLICATION SCHEDULE FOR TOMATO**

| DAYS AFTER PLANTATION | PRODUCTS | APPLICATION | METHOD AND DOSAGE |
|---|---|---|--|
| At the time of land preparation. | AJAY AZO, AJAY SPHURDA and AJAY POTASH, | To improve soil fertility naturally. | SOIL APPLICATION: Mix Ajay Azo- 2 Kg + Ajay Sphurda- 2 Kg + Ajay Potash- 2 Kg per acre . |
| 30 days after plantation | BIO-FIGHTER (EPN) | To preventive control of soil-borne larval pests in primary stage. | DRIP APPLICATION / SOIL APPLICATION- Biofighter - 1 Kg/Acre in 200 litres of water. |
| | AJAY VAM | For phosphorus solubilization, water absorption & root development. | SOIL APPLICATION: Mix 200 gm of AJAY VAM with 100 Kg of organic manure and then apply uniformly near the root zone. |
| | HUMIGUARD-PLUS | For white root development. | DRIP /DRENCHING APPLICATION- Humiguard Plus- 1 litre/Acre. |
| Two application- 30 days and 50 days after plantation. | GROWRICH PLUS | For healthy growth of the crop. | FOLIAR APPLICATION- Growth rich plus-2 ml/Litre of water. |
| | AGRONEEM 1% | To minimise infestation of sap sucking pests. | FOLIAR APPLICATION- Agroneem 1%- 1ml/litre of water. |
| | BIOSAN | To minimise infestation of bacterial and insect pest | FOLIAR APPLICATION- Biosan- 2 gm/litre of water. |
| | MITLAR | Biological management for sucking pests specially for Red Spider Mites. | FOLIAR APPLICATION- Mitlar-1ml/litre of water. |
| Three application- 30, 50 and 70 days after plantation. | KITOSTAR | Biological management for fungal and bacterial disease. | FOLIAR APPLICATION- Kitostar-2ml/litre of water. |
| | MICRO KOMBI-L | For balance nutrition to the crop. | DRIP / DRENCHING APPLICATION- Microkombi-L 500 ml in 200 Litres of water per acre. |
| | LIBRA | To improve flowering. | FOLIAR APPLICATION- Libra 0.5 to 1 ml/litre of water. |

- Repeat the sprays of dosages of Mitlar, Agroneem 1% and Kitostar, whenever required.
- * The above growth stages generally coincide with your plantation season. (State-wise plantation methods are different)
- ** The above given schedule taken as a Guideline for all stages of crop.
- Do not mix above Biological products with any chemical products. Spray during Early morning Or Late evening hours. Irrigation of Crops before 24-48 Hrs. of Bio-product applications is desirable.
- Keep minimum interval of 3 days between sprays of Biological products and Chemical products.

Note: The use of products of Ajay Biotech (India) Ltd. cannot be monitored by us in reality, we can only guarantee the quality of the products. Depending on the geographical area, spraying of fungicides and pesticides should be done according to and with the help of experts depending on crop disease, pest ratio, and climate change.